

Introduced marine species in Croatian waters (Eastern Adriatic Sea)

M. PEĆAREVIĆ¹, J. MIKUŠ¹, A. BRATOŠ CETINIĆ¹, J. DULČIĆ² and M. ČALIĆ³

¹University of Dubrovnik, Department of Aquaculture, Ćira Carića 4, 20000 Dubrovnik, Croatia

²Institute of Oceanography and Fisheries, Šetalište I. Meštrovića 63, 21000 Split, Croatia

³University of Dubrovnik, Institute for Marine and Coastal Research, Kneza Damjana Jude 12, 20000 Dubrovnik, Croatia

Corresponding author: marijana.pecarevic@unidu.hr

Handling Editor: Argyro Zenetos

Received: 24 October 2012; Accepted: 28 February 2013; Published on line: 26 March 2013

Abstract

The Croatian part of the Adriatic Sea covers more than 35% of the total Croatian territory, which means that monitoring changes in marine ecosystems and the conservation of biodiversity are of great importance. Following global changes, Croatia is experiencing increasing problems due to the introduction of new species that include aliens (due to aquaculture activities and shipping) and species from other Mediterranean subregions that are extending their geographic range. This work provides a checklist of introduced species in Croatian waters. A total of 113 species (15 phytoplankton, 16 zooplankton, 16 macroalgae, 44 zoobenthic and 22 fish species) have been recorded in the eastern part of the Adriatic Sea, of which 61 species are alien and 52 introduced, due to climate change.

Keywords: Alien species, range expansion, Croatia, Adriatic Sea, Mediterranean Sea.

Introduction

The Adriatic Sea is an elongated semi-enclosed basin in the Northeastern Mediterranean Sea. It extends in a southeast – northwest direction, with a shallow northern part and a deeper southern part. Adriatic Sea ecosystems are influenced by regular exchange of water with the Mediterranean Sea (Gačić *et al.*, 2010).

Water temperature in the deeper layers of the Adriatic Sea is above 11 - 12 °C during the whole year. In the warm part of the year, the seasonal thermocline is formed at 10 to 30 m of depth. In the summer, upper water layers reach temperatures of 22 - 25 °C (Buljan & Zore-Armanda, 1976). In the winter, the temperatures of open surface water fluctuates between 6 and 15 °C, with lower temperatures in the northern part and higher ones in the southern part of the basin (Pérès & Gamulin-Brida, 1973). During this period, vertical mixing in the water column occurs, thus transporting nutrients from the deep areas to the surface, which increases autotrophic production (Gačić *et al.*, 2002). The average salinity of the Adriatic Sea is 38.3 ‰, with the lower values observed during the winter months (Buljan & Zore-Armanda, 1976). The Adriatic Sea is characterized by varying trophic conditions; a productive and potentially eutrophic northern part, an oligotrophic southern part and also clearly oligotrophic open waters (Viličić *et al.*, 1989).

Considering all major characteristics, three biogeo-

graphic regions can be recognized in the Adriatic Sea: a shallow northern sub-basin and a deep southern sub-basin separated by a central sub-basin of intermediate depth (Zavatarelli *et al.*, 1998).

The first studies of Adriatic Sea flora and fauna started in the early 18th century and a comparatively large amount of data has been collected till the present time (Zavatarelli *et al.*, 1998).

Like many other countries, Croatia is facing increasing problems due to the introduction of new species. Although several projects on the biological and ecological characteristics of the Eastern Adriatic ecosystems are being conducted continuously and the results of such research is published in scientific literature, there is still no national data base or report bringing together all available data on introduced species in the Adriatic Sea. Therefore, the aim of this paper is to provide complete data about introduced marine species in Croatian seawaters, whether introduced by Man (alien) or unintentionally due to climate change.

Materials and Methods

This article comprises all available data regarding introduced species in the Croatian part of the Adriatic Sea (the Eastern Adriatic Sea). It brings together data from available scientific literature, reports, websites such

as MAMIAS (UNEP-MAPRAC/SPA, 2013) as well as personal knowledge and unpublished data. Phytoplankton species designated as new in Croatian Sea are those that have not been listed in the most recent checklist of Eastern Adriatic phytoplankton (Viličić *et al.*, 2002). The status of introduced species for other taxa was assigned to species of foreign origin, which were recorded in the Adriatic Sea. Certain species were rejected as non indigenous species, since new knowledge demonstrated that they should be considered as native species.

A list of all introduced species in the Eastern Adriatic Sea is presented in two tables. Table 1 includes non indigenous species whose findings are likely due to maritime transport, aquaculture or an unknown vector but classified as “aliens” in Zenetos *et al.* (2010), while Table 2 comprises vagrant species of Atlantic and Indo-Pacific origin and Mediterranean species which have entered the Eastern Adriatic Sea recently, presumably due to climate change.

The taxonomy of the species presented in this paper follows the WoRMS (World Register of Marine Species) (Appeltans *et al.*, 2012).

Results and Discussion

Introduced planktonic species in the Eastern Adriatic Sea

Introduced phytoplankton

Akashiwo sanguinea (K. Hirasaka) G.Hansen & Ø.Moestrup, 2000 is often found in samples from the Southern and Middle Eastern Adriatic (Carić *et al.*, 2011), but is not included in the Eastern Adriatic phytoplankton checklists (Viličić *et al.*, 2002); it was, therefore, regarded as a new species in Croatian waters.

Ceratoperidinium yeye Margalef, 1969 was found in net samples from the coastal waters of the Eastern Adriatic Sea in both the Northern and the Middle Adriatic during the summer of 2003, when seawater temperature was above 20 °C (Ninčević Gladan *et al.*, 2006). *Ceratoperidinium* is a rare dinoflagellate genus reported from both the western and the eastern Mediterranean Sea (Margalef, 1969; Gómez & Abboud-Abi, 2003). Literature data on *Ceratoperidinium* ecology and distribution is very scarce.

Nine new species for the Croatian phytoplankton database, *Gymnodinium abbreviatum* Kofoid et Swezy, 1921, *G. agilisforme* Schiller, 1928, *G. grammaticum* (Pouchet) Kofoid et Swezy, 1921, *G. ostenfeldii* Schiller, 1928, *G. uberrimum* (G.J.Allman) Kofoid & Swezy, 1921, *Gyrodinium ascendans* Kofoid, 1931, *G. obtusum* (Schütt) Kofoid & Swezy, 1921, *G. pingue* (Schütt) Kofoid & Swezy, 1921 and *Woloszynskia neglecta* (Schilling) R.H.Thompson, 1950, were recorded at the Maslinova Bay fish farm (Skejić *et al.* 2012). All these species are cosmopolitan and thus their origin in the Adriatic remains unknown.

Neoceratium paradoxides (Cleve) F.Gomez, D.Moreira & P.Lopez-Garcia, 2009 and *Ornithocercus splendidus* F.Schütt, 1895 are two newly recorded phytoplankton species in the Adriatic Sea, found for the first time in the deep South Adriatic in 2008 (Batistić *et al.*, 2012). Both are known from the northern Levantine Basin (Polat & Koray, 2007) so their occurrence might be related to documented changes in Ionian Sea circulation that result in the flow of Aegean and Levantine waters into the Adriatic Sea (Batistić *et al.*, 2012).

Ostreopsis ovata Fukuyo, 1981 is an epiphytic, potentially toxic dinoflagellate living on macroalgae, found in the Northern Adriatic for the first time in 2006 (Monti *et al.*, 2007). *O. ovata* was isolated from macroalgae in two areas, the Gulf of Trieste (Italy) and close to Rovinj (Croatia) and was identified by scanning electron microscopy and by fluorescence light microscopy.

Scaphodinium mirabile Margalef, 1963, often reported in the Mediterranean Sea (Gómez & Gorsky, 2003), was recorded for the first time in the Eastern Adriatic Sea in Mali Ston Bay in 2002, in the shellfish farm area (Čalić, 2010) and later, in 2011, was also found in the Port of Ploče.

Introduced zooplankton

Arctapodema australis (Vanhöffen, 1902) and *Eucheilota paradoxica* Mayer, 1900 were found in samples from the Southern Adriatic collected in 1967 (Schmidt & Benović, 1977). *A. australis* was rare until 2002 but after that it was frequently found in Croatian waters and today it represents one of the typical Southern Adriatic medusae (Lučić *et al.*, 2009). The ability of hydroids to produce resting stages and settle on floating objects (including ships) enables them to achieve a wide distribution (Cornelius, 1981).

Arietellus pavoninus Sars G.O., 1905 is a tropical copepod which was presumably ship transferred (e.g. ballast water) via the Suez Canal or via the Gibraltar Strait (Kovalev, 2006) and was found in the Eastern Adriatic Sea (Hure & Kršinić, 1998).

Corymorpha annulata (Kramp, 1928) was reported in the Adriatic Sea (Schmidt & Benović, 1977), but since a single specimen was found with no additional records, more observations are needed to confirm the presence of this rare Indo-Pacific medusa in the Adriatic Sea.

The pteropod *Desmopterus papilio* Chun, 1889 and the heteropod *Protatlanta souleyeti* (E. A. Smith, 1888) were found for the first time in the Adriatic Sea during investigation at a deep-sea station in the northern part of the South Adriatic Pit (Batistić *et al.*, 2004).

Haliscera bigelowi Kramp, 1947 is a typically bathypelagic hydromedusa recorded for the first time in the Adriatic Sea in spring 2002 at a deep-sea station in the South Adriatic (Benović *et al.*, 2005). It is probably a recent invader of the Adriatic, as are certain other planktonic cnidarians (Kršinić & Njire, 2001).

The presence of the immigrating Atlantic calycophoran *Lensia subtiloides* (Lens & van Riemsdijk, 1908) was recorded in the Southern Adriatic Sea (Batistić & Garić, 2012).

The calycophoran siphonophore *Muggiaea atlantica* Cunningham, 1892 was recorded for the first time in the coastal southeastern (Gamulin & Kršinić, 2000) and central Adriatic (Batistic, 1999). After these first records it has been observed that native species *Muggiaea kochi* is progressively being replaced by *M. atlantica* in the coastal waters of the Eastern Adriatic, especially in eutrophicated areas (Batistić *et al.*, 2007).

Niobia dendrotentaculata Mayer, 1900, a tropical Atlantic origin species, after its first record in the Adriatic in 2001, again occurred in August 2006 and accounted for 44% of total hydromedusan abundance (Batistić & Garić, 2010).

Peniculus fistula Von Nordmann, 1832 is a parasitic pennellid copepod, wide spread in the Atlantic and Indo-Pacific waters, and relatively common in Mediterranean fishes (Raibaut *et al.*, 1998). This species was recorded for the first time in the Eastern Adriatic Sea during an investigation of the biology of garfish *Belone belone* Linnaeus, 1761 (Vidjak *et al.*, 2008). Garfish were caught from February to March 2008 near the Islands of Korčula and Dugi Otok.

Polychaetes *Phalacrophorus pictus* Greef, 1879 and *Pontodora pelagica* Greef, 1879 were found in plankton samples from the Southern Adriatic Sea during the investigated period 1993-2011 (Batistić & Garić, 2012).

An occurrence of *Sagitta galerita* Dallot, 1971 was recorded in the Croatian waters of the Southern Adriatic (Batistić & Garić, 2012).

Thalia orientalis Tokioka, 1937 has been found in Croatian waters since January 2008 and has replaced its formerly dominant congener *T. democratica* in the coastal and open Southern Adriatic waters (Batistić *et al.*, 2009).

Trichydra pudica Wright, 1858 was recorded in the Eastern Adriatic Sea for the first time in 1973 (Schmidt & Benović, 1979).

Introduced benthic species in the Eastern Adriatic Sea *Introduced phytobenthos*

Acrothamnion preissii (Sonder) E.M. Wollaston, 1968 is a tropical filamentous alga which was first recorded in the Eastern Adriatic in 2007 when it was found near Dubrovnik (Despalatović *et al.*, 2008). This invasive red alga has a negative impact on domestic benthic and epiphytic communities because it creates turfs, which outgrow native species of algae and prevent their settlement (Despalatović *et al.*, 2008).

Asparagopsis armata Harvey, 1855 was introduced in the Mediterranean Sea unintentionally with oysters

(Ribera & Boudouresque, 1995) and naturally through the Strait of Gibraltar (Pacios *et al.*, 2011); it is widely distributed along the Western Mediterranean coastline. In the Adriatic Sea it was first reported in 1978 in Montenegro (Špan & Antolić, 1983) and later in 1991 in the Gulf of Trieste (Giaccone, 1978). In the same year, the tetrasporophyte (*Falkenbergia* phase) of this alga was found in Slovenian coastal waters and in 1997 gametophyte plants were found in Croatian waters, near Senj (Orlando-Bonaca, 2010).

Asparagopsis taxiformis (Delile) Trevisan de Saint-Léon, 1845 is a red alga whose gametophyte phase is spreading along Mediterranean coasts carried by currents or on ship hulls (Boudouresque & Verlaque, 2002). The first record in the Eastern Adriatic Sea is from 2007 when specimens of *A. taxiformis* were found on rocky shallow water substrate in the Dubrovnik area, and also in the Mljet National Park (Despalatović *et al.*, 2008).

Caulerpa racemosa var. *cylindracea* (Sonder) Verlaque, Huisman & Boudouresque, 2003 is a tropical green alga aggressively widespread throughout the Mediterranean basin since the early 1990s (Piazzi *et al.* 2001). It was found for the first time in Croatia in autumn 2000 near the Pakleni Islands (Žuljević *et al.*, 2003), and has now been recorded along the entire coast, mostly in the Southern and Middle Adriatic Sea (Žuljević *et al.*, 2010).

Caulerpa taxifolia (Vahl) Agardh, 1817 is a tropical green alga whose first occurrence in the Mediterranean was recorded in Monaco in 1984; since then it has been successfully invading Mediterranean coasts (Meinesz *et al.*, 2001). In the Eastern Adriatic, the species was initially observed at three locations: Island of Hvar and Island of Krk in 1994, and Barbat Channel in 1996 (Špan *et al.*, 1998). The records from Malinska and the Harbor of Vrstar represent the highest northern latitudes at which this invasive alga has been found (Iveša *et al.*, 2006).

Chondria coerulescens (J. Agardh) Falkenberg, 1901 was recorded in Rijeka Bay (Battelli & Arko Pijevac, 2003) where this red alga has established a population (Antolić *et al.*, 2011a).

Chondria pygmaea Garbary & Vandermeulen, 1990 is an epiphytic red alga found for the first time along the Croatian coast, in the Gulf of Kvarner, in 1997 (Sartoni & Rossi, 1998). This finding is important because it is the most western occurrence of this alga and shows that *C. pygmaea* is a successful migrant, which can develop within a wide ecological range (Sartoni & Rossi, 1998).

Codium fragile ssp. *fragile* (Suringar) Hariot, 1889 is a green alga which is widespread thanks to its high environmental tolerance and has also extended its range to the Adriatic Sea (UNEP-MAP-RAC/SPA, 2013).

Colpomenia peregrina Sauvageau, 1927 is called the "oyster thief" because it attaches to oysters and then floats off with them. Records from the Eastern Adriatic are rare, but its presence there is possible (Cabioch *et al.*, 1992).

Desmarestia viridis (O.F.Müller) J.V.Lamouroux, 1813 has only four records in the Mediterranean Sea, and one of them is from the Eastern Adriatic coast where this alga was found in 1948 (Occhipinti Ambrogi, 2002). However, according to Boudouresque and Verlaque (2002) some populations may be native in the Northern Adriatic.

Lophocladia lallemandii (Montagne) F. Schmitz 1893 is an invasive macroalga recorded for the first time in the Eastern Adriatic Sea in 1970 (UNEP-MAP-RAC/SPA, 2013).

Parviphycus antipai (Celan) B.Santelices, 2004 has been recorded frequently in the Mediterranean Sea (Boudouresque, 1972; Boisset, 1986) and probably overlooked in the Adriatic Sea. The first record from the Adriatic Sea was in 1997 when a few isolated plants, growing in epiphytic communities on crustose fleshy species, were collected near the Island of Cres (Sartoni & Rossi, 1998).

Polysiphonia banyulensis Coppejans, 1976 has been recorded frequently in the North-Western Mediterranean (Cossu *et al.*, 1993). The 1997 record from the Adriatic Sea extends the distribution area of this epiphytic red alga, which is now known from widely separated geographical areas (Sartoni & Rossi, 1998).

Polysiphonia paniculata Montagne, 1842 was found in Croatian waters in 1975 (UNEP-MAP-RAC/SPA, 2013). This red alga was also included in the recent check list of the benthic macroflora from Rijeka Bay (Antolčić *et al.*, 2011b).

Ulva fasciata (= *U. lactuca sensu Linnaeus*) was recorded in the middle part of the Eastern Adriatic Sea (Ercegović, 1980).

Womersleyella setacea (Hollenberg) R.E.Norris, 1992 is a tropical alga that was first noted in the Adriatic Sea in 1997, near the Island of Cres (Sartoni & Rossi, 1998). Later on it was found near Cape Oštro and Rabac (Batelli & Arko-Pijevac, 2005) and near the Island of Dugi Otok (Kružić, 2008b). Nikolić *et al.* (2010) assume that the entire coastline of the Adriatic Sea is affected by this invasive alga.

Introduced zoobenthos

Two barnacle species, namely *Amphibalanus eburneus* (Gould, 1841) and *A. improvisus* (Darwin, 1854), were recorded in the Rovinj area (Zavodnik & Igić, 1968; Igić, 1986) where they were probably introduced by shipping activity (Galil *et al.*, 2011).

Anadara transversa (Say, 1822) is a bivalve that was recently recorded as empty complete shells from the Adriatic Croatian shores (Peharda *et al.*, 2010). Six specimens of *A. transversa* were found in Lim Bay, Northern Adriatic, attached by byssus to a tile near a mesh bag of *Mytilus galloprovincialis* at a depth of 4.4 m (Nerlović *et al.*, 2012).

Aplysia dactylomela Rang, 1828 is a gastropod firstly recorded in the Eastern Adriatic near the Island of Sušac

in the 2006 (Turk, 2006). It appears that this species also lives in the Mljet area (Despalatović *et al.*, 2008).

Arcuatula senhousia (Benson in Cantor, 1842) is a Western Pacific bivalve, found in Savudrija Bay in 2003 by local fisherman (Crocetta, 2011).

Aspidosiphon (Akrikos) mexicanus (Murina, 1967) is an Atlantic sipunculan species recorded in the Eastern Adriatic in 1986 (Murina & Zavodnik, 1986) and its present status in Croatian waters is unknown (UNEP-MAP-RAC/SPA, 2013).

Astroides calycularis (Pallas, 1767) is a warm water coral which was recorded for the first time in the Adriatic Sea in 1899 in Rijeka Bay, but there are many doubts about correct determination (Zibrowius & Grieshaber, 1977; Zavodnik & Kovačić, 2000). During the last century it was recorded along the entire Eastern Adriatic coast (Pax & Müller, 1962; Grubelić *et al.*, 2004). The recent northernmost record is from the rocky shore of the Islet of Borovik in the Northern Adriatic (Kružić *et al.*, 2002).

Balanus trigonus Darwin 1854 is an introduced barnacle, which represents an example of anthropogenous cosmopolitanism in the Adriatic Sea (Igić, 2007). It was probably introduced in the Adriatic Sea by fishing ships from the northwestern coast of Africa (Relini, 1968).

Brachidontes pharaonis (P. Fischer, 1870) is an Indo-Pacific mytilid with a highly variable shell and is considered to be a variant of *Brachidontes ustulatus* (Lamarck, 1819) (Huber, 2010). It was firstly mentioned in the Croatian Adriatic by Hrs-Brenko and Legac (2006). Orlando Bonaca (2001) listed it as "occasional" and no follow-up is known from the area (Despalatović *et al.*, 2008). The findings in Croatia are likely due to maritime transport linked with the oil terminal of Trieste.

Bugula fulva Ryland, 1960 is an Atlantic bryozoan found in samples taken along and just offshore of the Istrian Peninsula and Rovinj area (Hayward & McKinney, 2002).

Bursatella leachii De Blainville, 1817 is a gastropod with a worldwide distribution in warm temperate and tropical waters. This Lessepsian migrant was recorded for the first time in the Eastern Adriatic in the Rovinj area (De Min & Vio, 1998). After that it was also reported from the Island of Hvar and the Split area (Despalatović *et al.*, 2008). *Callinectes sapidus* was firstly found in the Eastern Adriatic Sea in 2004 near Ston in a hypersaline lagoon (salt ponds) (Onofri *et al.*, 2007). Findings in the Neretva River estuary in 2004 and 2006 confirm the spreading of this species throughout the Adriatic Sea (Dulčić *et al.*, 2011b).

Cladocora debilis Milne Edwards & Haime, 1849, a scleractinian coral, was found for the first time in spring 2002 on the cliffs, south of Lastovo Island (Kružić *et al.*, 2007).

Cladopsammia rolandi Lacaze-Duthiers, 1897 is a colonial scleractinian coral which was recorded for the first time in the Adriatic Sea in 2002, when colonies were found in the Lastovo Island area (Kružić, 2008a). In 2005, numerous colonies were found at two locations, in the

south-western part of the Island of Mljet (Kružić, 2008a). Those findings in the Southern Adriatic are the northernmost findings of *C. rolandi* in the Mediterranean Sea.

Eight coral species, *Alicia mirabilis* Johnson, 1861, *Balanophyllia* (*Balanophyllia*) *regia* Gosse, 1853, *Caryophyllia* (*Caryophyllia*) *cyathus* (Ellis & Solander, 1786), *Coenocyathus cylindricus* Milne Edwards and Haime, 1848, *Dendrophyllia ramea* (Linnaeus, 1758), *Guynia anulata* Duncan, 1872, *Halcampoides purpureus* (Studer, 1879) and *Sphenotrochus* (*Sphenotrochus*) *andrewianus* Milne Edwards and Haime, 1848 were recorded for the first time in the Adriatic Sea during the “Thais” expedition, in the period 1995-1998 (Kružić, 2002). Since these findings do not indicate recent settlements it is possible that limited populations were previously overlooked (Kružić, 2002).

Coscinospira hemprichii Ehrenberg, 1839, *Cushmanina striatopunctata* (Parker & Jones, 1865), *Elphidium striatopunctatum* (Fichtel & Moll, 1798) and *Planispirinella exigua* (Brady, 1879) are introduced benthic foraminifera recorded in Croatian waters (UNEP-MAP-RAC/SPA, 2013). *C. hemprichii* was also found later in the lakes of Mljet Island (Vaniček *et al.*, 2000).

Crassostrea gigas (Thunberg, 1793) larvae were recorded in Limski Bay in the Northern Adriatic Sea (Hrs-Brenko, 1982). *C. gigas* settles on boats and ships and is often found in mariculture areas; therefore, anthropogenic activity might be the vector of introduction.

Eudendrium carneum Clarke, 1882 was perhaps introduced in the Mediterranean by human activity, as it often occurs on ship hulls and is thus potentially easily introduced to other regions (Millard, 1975); it has also been found in the Eastern Adriatic (UNEP-MAP-RAC/SPA, 2013). It is possible that besides *E. carneum*, another species from this genus, namely *E. cf. merulum*, Watson, 1985 is also present in Croatian waters (UNEP-MAP-RAC/SPA, 2013).

Ficopomatus enigmaticus (Fauvel, 1923) is a cryptic species; the first Adriatic record is from the Venetian lagoons in 1934 and today it builds huge aggregate “reefs” in the Po River Delta (Bianchi & Morri, 1996). The species is also established in Slovenia (Lipej *et al.*, 2012). In Croatian waters, it was recorded for the first time in 2006 in the Bay of Šibenik (Mikuš *et al.*, 2007). Recent records from two new locations along the eastern coast of the Adriatic Sea, Krka River Estuary and Neretva River Delta, have been reported in 2009 (Cukrov *et al.*, 2010).

Halgerda willeyi, Eliot 1903 is a nudibranch recorded for the first time in Croatian waters in July 1988, at the southernmost tip of the island of Cres, near Baldarin Bay (Turk, 2000).

Hemigrapsus sanguineus (de Haan, 1835) was recorded in August 2001 along the northwestern coast of the Peninsula of Istria, when a single adult male specimen of this East Asian crab was collected (Schubart, 2003). The current absence of *H. sanguineus* in southern Europe and the western Mediterranean suggests an in-

dependent human-mediated introduction of the Croatian specimen (Schubart, 2003).

Hippolyte prideauxiana Leach, 1817 is a caridean shrimp, collected in 2002 at Kostrena, near the city of Rijeka, in the Northern Adriatic. This record, the first for the Adriatic Sea, widens the previously known geographic distribution of this species (Kirinčić, 2006).

Idas simpsoni (J.T. Marshall, 1900) is a rare bivalve mollusc that attaches itself by byssus to the skull of the whales. It was recorded for the first time in the Adriatic Sea in 2003 on a skull that was trawled up from a depth of 430 m near the Island of Mljet. On that occasion, more than 127 specimens of *I. simpsoni* were collected from the skull of a fin whale *Balaenoptera physalus* (Linnaeus, 1758) (Bolotin *et al.*, 2005).

Marteilia refringens Grizel, Comps, Bonami, Cousserans, Duthoit & Le Pennec, 1974 is a protistan parasite responsible for marteiliosis in the flat oyster *Ostrea edulis* and the mussels *Mytilus edulis* and *M. galloprovincialis* respectively. Zrnčić *et al.* (2001) detected *Marteilia* sp. in *Mytilus galloprovincialis* in Croatia.

Megabalanus tintinnabulum (Linnaeus, 1758) was found near Rovinj, Pula and Rijeka, reported as *Balanus tintinnabulum* (Kolosváry, 1947).

Melibe viridis (Kelaart, 1858) is an Indo-Pacific gastropod recorded along all Mediterranean coasts, including Croatian, where it was found for the first time in 2001 near the Island of Hvar, which is the northernmost record in the Mediterranean (Despalatović *et al.*, 2002).

Metasychis gotoi (Izuka, 1902) and *Nereis persica* Fauvel, 1911 were reported as introduced species in Croatian waters but their status is questionable (UNEP-MAP-RAC/SPA, 2013).

Neopseudocapitella brasiliensis Rullier & Amoureux, 1979 was found in the harbour and the near vicinity of Rovinj (Zavodnik *et al.*, 1985).

Ochetostoma erythrogrammon Leuckart & Ruppell, 1828 was reported for the first time in 1962 when a sample from Velebit Canal was found to contain four specimens. This is the first report of the genus from the Mediterranean Sea (Saiz Salinas & Ruthensteiner, 2005).

Phascolosoma scolops (Selenka & de Man, 1883) was found in the northern part of the Croatian Adriatic Sea (Murina, 1976) and is the only one found in the Mediterranean (Murina & Zavodnik, 1986).

Pinctada imbricata radiata (Leach, 1814) is a species known as a pearl oyster, recorded for the first time in Croatian waters in 2006, when two juvenile specimens were found in the northern part of the Adriatic Sea coast, off Pula, at a depth of 59 m (Dogan & Nerlović, 2008). The presence of *P. imbricata radiata* in this area could be the result of shipping since this finding is near the location of previous findings in the Trieste Bay (Dogan & Nerlović, 2008).

Siphonaria pectinata (Linnaeus, 1758) is a warm water gastropod, recorded for the first time in the Eastern

Adriatic in 2003, when a numerous population was found in the Split area (Despalatović *et al.*, 2008). The possible vector of introduction is shipping, as two international maritime transport harbours are located in the vicinity of the location of the first record.

Introduced ichthyofauna in the Eastern Adriatic Sea

Alectis alexandrinus (Geoffroy Saint-Hilaire, 1817) is a species that is distributed in the southern parts of the Mediterranean Sea. One specimen, caught near Split in summer 1973 and kept in the Natural Museum in Genova (Italy), was scientifically proven as *A. alexandrinus* in 2005 (Dulčić, 2005).

Caranx crysos (Mitchill, 1815) is an Atlantic species, recorded for the first time in the Adriatic Sea near Plićina Ćrvar (western coast of the Peninsula Istria) in 2008 (Dulčić *et al.*, 2009b). This is the northernmost finding of this species in the Mediterranean. There are no additional records for this species in the Adriatic Sea.

Caranx rhonchus Geoffroy Saint-Hilaire, 1817 is a common predatory fish in the eastern Atlantic and in 2011 it was recorded for the first time in the southern part of the Eastern Adriatic Sea in Mali Ston Bay (Kožul & Antolović, 2013).

Cyclopterus lumpus Linnaeus, 1758 naturally occurs in the Atlantic, and it was found in the Adriatic Sea for the first time near Molunat Bay in 2004 (Dulčić & Golani, 2006). Shipping, e.g. ballast water, is the most probable vector of introduction. There are no additional records for this species in the Adriatic Sea.

Elates ransonnetii (Steindachner, 1876) normally lives in the Pacific Ocean, from south Chinese shores to north Australia. The first record in the Adriatic Sea is from 2010, when a single specimen was caught in Kaštela Bay (Dulčić *et al.*, 2010). There are no additional records for this species in the Adriatic Sea.

Enchelycore anatina (Lowe, 1838) is a subtropical species that was recorded for the first time in the Adriatic Sea in 2010. Two specimens were recorded using an underwater camera near the Island of Sušac (Lipej *et al.*, 2011). There are no additional records for this species in the Adriatic Sea.

Epinephelus aeneus (Geoffroy Saint-Hilaire, 1817) is a thermophilic species from the eastern Atlantic and the southern Mediterranean Sea. This species was recorded for the first time in 1998 near Dubrovnik (Glamuzina *et al.*, 2000) and additionally near the Island of Dugi Otok in 2006 (Dulčić *et al.*, 2006). There is evidence of northward spreading of groupers, and this is the northernmost occurrence of this species.

Equulites (Leiognathus) klunzingeri (Steindachner, 1898) is a Lessepsian migrant, recorded in the Adriatic Sea for the first time near the Island of Mljet in 2000 (Dulčić & Pallaoro, 2002). There are no additional records for this species in the Adriatic Sea.

Fistularia commersonii Rüppel, 1838 is a Lessepsian migrant, recorded in the Adriatic Sea for the first time in 2006 near the Island of Sv. Andrija and near the Italian coast (Bari) (Dulčić *et al.*, 2008). There are also new additional records in Montenegro (Joksimović *et al.*, 2008) and the Island of Šipan (two juvenile specimens). Based on additional records, the existence of a self-sustaining population in the Adriatic is possible.

Holacanthus ciliaris (Linnaeus, 1758) was found for the first time in the Adriatic Sea, Trogir Bay, in 2011. So far, this is the only record in the Eastern Adriatic Sea.

Lagocephalus lagocephalus lagocephalus (Linnaeus, 1758) is a thermophilic species distributed in tropical and subtropical waters in the Atlantic, Indian and Pacific Oceans. The first report of the species is from the Southern Adriatic, from the area near Molunat Bay in 2004 (Dulčić & Pallaoro, 2006). There are no additional records for this species in the Adriatic Sea.

Lobotes surinamensis (Bloch, 1790) is a tropical and subtropical species. It was recorded for the first time in the Adriatic Sea in 2010 near the Island of Biševo (Dulčić & Dragičević, 2011a). Based on the additional records (Island of Šipan), the presence of a self-sustaining population in the Adriatic is possible.

Mycteroperca rubra (Bloch, 1793) is a thermophilic species distributed in the eastern Atlantic and Mediterranean Sea. This species was recorded for the first time in 2000 near Dubrovnik (Glamuzina *et al.*, 2002). There are some reports of additional records of this species in the Adriatic.

Pagrus major (Temminck & Schlegel, 1843) was recorded for the first time in the Adriatic Sea in the Zadar archipelago, near the Island of Molat, in 2004 (Dulčić & Kraljević, 2007). Its presence in the Adriatic is probably due to the escape from mariculture facilities. Based on the additional records, the existence of a self-sustaining population in the Adriatic is possible.

Pampus argenteus (Euphrasen, 1788) is an Indo-Pacific species; the only specimen found in the Mediterranean Sea was caught in the Eastern Adriatic Sea, near Rijeka, in 1896. This was the first Lessepsian migrant reported in the Mediterranean Sea (Dulčić *et al.*, 2004). Besides active migration as a probable vector of introduction, another possible vector of introduction could be shipping.

Paranthias furcifer (Valenciennes, 1828) was found in Marina Bay near Trogir in 2011 (Pećarević & Mikuš, 2012). This fish was found near an oil platform that arrived from the Gulf of Mexico and, therefore, shipping could be the vector of introduction.

Plectorhinchus mediterraneus (Guichenot, 1850) is a thermophilic species distributed along the West African coast and in the Western Mediterranean. The first records for the Adriatic coast were reported in 1993 from Trieste Bay and Savudrija Bay (Lipej *et al.*, 1996). There are no additional records for this species in the Adriatic Sea.

Siganus luridus (Rüppel, 1829) is a Lessepsian mi-

grant, recorded in the Adriatic Sea for the first time in the summer of 2010 in Trieste Bay (Poloniato *et al.*, 2010). There are some additional records of this species from the Island of Mljet in 2010 (Dulčić *et al.*, 2011a) and the Island of Šipán in 2012, and it seems that this species has established a population in the Adriatic Sea.

Siganus rivulatus Forsskål, 1775 is a Lessepsian migrant, recorded in the Adriatic Sea for the first time in 2000 near the Island of Bobara in the Southern Adriatic (Dulčić & Pallaoro, 2004). There are some additional records of this species that have not been verified yet.

Sphoeroides pachygaster (Müller & Troschel, 1848) was recorded in Croatian waters for the first time in 1992 (Pallaoro & Jardas, 1996). A juvenile specimen was captured again in 2002, off Cape Kamenjak (Peninsula of Istria, Northern Adriatic) and represents the northernmost report in the Adriatic Sea and the entire Mediterranean area (Dulčić, 2002).

Sphyraena chrysotaenia Klunzinger, 1884 is a Lessepsian migrant, recorded in the Adriatic Sea for the first time in 2000 near Molunat Bay in the Southern Adriatic (Pallaoro & Dulčić, 2001). According to several additional records, this species has established a population.

Sphyraena viridensis Cuvier, 1829 is a thermophilic species distributed in the eastern Atlantic and the Mediterranean Sea. The first record in the Adriatic Sea from the area of Dubrovnik in the Southern Adriatic in 2003 (Kožul *et al.*, 2005), was followed by records from the Northern Adriatic, near the Port of Rijeka in 2008 (Dulčić *et al.*, 2009a). Several specimens were caught in the Middle Adriatic as well, and it seems that this species has established a population in the Adriatic Sea.

In total, at least 113 introduced species have been recorded in the eastern part of the Adriatic Sea, including 15 phytoplankton, 16 zooplankton, 16 macroalgae, 44 zoobenthic and 22 fish species (Tables 1 and 2).

For the majority of the introduced phytoplankton species, the vector of introduction might be related to climate change, areal expansion and changes in sea circulation resulting in the flow of Aegean and Levantine waters into the Adriatic Sea. *Gymnodinium* is a relatively little-known genus with poorly described species on the eastern coast of the Adriatic Sea, Viličić *et al.* (2002) recorded three species in the entire eastern basin.

Sixteen introduced zooplankton species are listed in this paper. However, some of the protozoan planktonic species, reported as non indigenous species for the Northern Adriatic, i.e. tintinnid ciliates *Tintinnopsis mortenseni* Schmidt, 1901 (found in the ballast water of a ship in the Port of Koper) and *Codonellopsis orthoceras* Haeckel (Jørgensen), 1924, have already been found in other parts of the Adriatic Sea (David *et al.*, 2007). *T. mortenseni* usually occurs in the Middle Adriatic Sea (Kaštela Bay) in March (Kršinić, 1980), while *C. orthoceras* was found

in the surface layer of the open waters of the Southern Adriatic (Kršinić & Grbec, 2006). That is the reason why those species were not considered as non indigenous for the Eastern Adriatic.

Introduced benthic algae were found along the entire coast of the Eastern Adriatic. Fast establishment of their populations and successful expansion were enabled by the lack of natural predators and/or high possibility of vegetative reproduction. In favourable new environments these species have become invasive and have a negative impact on natural habitats and communities. Introduced green algae in the Adriatic Sea give rise to major concerns since two non indigenous *Caulerpa* species have rapidly spread across a large part of the Croatian coast overthrowing native benthic communities (Žuljević *et al.*, 2010). There are certain indications that the secondary spread of both *Caulerpa* species has been caused by shipping activities, e.g. ballast waters, anchors, fishing nets, etc. (Verlaque *et al.*, 2003). Introduced red algae were reported in the Eastern Adriatic Sea from the 1997. For some of them the invasive character has been well documented (e.g. *Womersleyella setacea* (Hollenberg) R.E.Norris, 1992). Streftaris and Zenetos (2006) classified *W. setacea* among the 100 worst marine alien species in the Mediterranean, and there is a great chance that this species could be assigned the status of the worst alien macrophyte in the Adriatic Sea as well. New records of brown algae are not so frequent in the Eastern Adriatic, but there are two potentiality introduced species. Most of the vectors of introduction of benthic algae are unknown, but the human factor, e.g. shipping and aquaculture, is not excluded.

Introduced zoobenthic species are quite numerous in the Eastern Adriatic. Introduction of some zoobenthic species is related to climate change and range expansion, while for other species the vectors of introduction are mainly associated with shipping activities. Recent research on the zoobenthos in Croatian waters has resulted in the discovery of numerous scleractinian species that proved to be new to the area (Kružić, 2002) and the previously uncertain occurrence of certain species (e.g. *A. calycularis* and *D. ramea*) was confirmed (Kružić *et al.*, 2002).

A large number of a new fish species has been recorded in the last few decades, and the abundance of species that were previously rare in the Adriatic Sea has also increased. The highest growth was observed among thermophilic species, which were firstly present in the Southern Adriatic but can now be found at more northern latitudes too. Lessepsian migrant *Stephanolepis diaspros* Fraser-Brunner, 1940 and thermophilic species *Tylosurus acus imperialis* (Rafinesque, 1810) were found far south, off the Croatian coast. The findings were reported for the Montenegro coast (Bello, 1995; Dulčić & Pallaoro, 2003). Additional records suggest that *T. acus imperialis* has established a population in the Adriatic Sea. However,

Table 1: List of non indigenous species introduced in the Eastern Adriatic Sea (Croatia) by human or unknown vectors. Legend: Location: NA – Northern Adriatic, MA – Middle Adriatic, SA – Southern Adriatic; Vector: AQ – aquaculture, LM – Lessepsian migrant, SH – shipping, , UN – unknown; Status: C – casual, E – established, S – spreading.

Taxon	First record	Location	Vector	Status	Reference
Pyrrhophyta (Dinophyceae)					
<i>Ostreopsis ovata</i>	2006	NA	UN	C	Monti <i>et al.</i> , 2007
Chlorophyta					
<i>Caulerpa racemosa</i>	2000	MA	SH	E	Žuljević <i>et al.</i> , 2003
<i>Caulerpa taxifolia</i>	1994	MA	SH	E	Špan <i>et al.</i> , 1998
<i>Codium fragile</i> ssp. <i>fragile</i>	1983	NA	UN	E	UNEP-MAP-RAC/SPA, 2013
<i>Ulva fasciata</i>	<1969	MA	UN	C	Ercegović, 1980
Rhodophyta					
<i>Acrothamnion preissii</i>	2007	SA	UN	C	Despalatović <i>et al.</i> , 2008
<i>Asparagopsis armata</i>	1997	NA	UN	S	Orlando-Bonaca, 2010
<i>Asparagopsis taxiformis</i>	2007	SA	UN	C	Despalatović <i>et al.</i> , 2008
<i>Chondria coerulescens</i>	1997	NA	UN	E	Battelli & Arko Pijevac, 2003
<i>Chondria pygmaea</i>	1997	NA	LM	C	Sartoni & Rossi, 1998
<i>Lophocladia lallemandii</i>	1970	NA	UN	E	UNEP-MAP-RAC/SPA, 2013
<i>Polysiphonia paniculata</i>	1975	NA	UN	C	UNEP-MAP-RAC/SPA, 2013
<i>Womersleyella setacea</i>	1997	NA	UN	E	Sartoni & Rossi, 1998
Pheophyta					
<i>Colpomenia peregrina</i>	<1992	-	UN	C	Cabioč'h <i>et al.</i> , 1992
<i>Desmarestia viridis</i>	1948	NA	UN	C	Occhipinti Ambrogio, 2002
Foraminifera					
<i>Coscinospira hemprichii</i>	1911	NA, SA	UN	S	UNEP-MAP-RAC/SPA, 2013
<i>Cushmanina striatopunctata</i>	1913	NA	UN	C	UNEP-MAP-RAC/SPA, 2013
<i>Elphidium striatopunctatum</i>	1911	NA	UN	C	UNEP-MAP-RAC/SPA, 2013
<i>Planispirinella exigua</i>	1910	NA	UN	C	UNEP-MAP-RAC/SPA, 2013
Cercozoa					
<i>Marteilia refringens</i>	1998	NA	UN	C	Zrnčić <i>et al.</i> , 2001
Cnidaria					
Hydrozoa					
<i>Arctapodema australis</i>	1967	SA	SH	E	Schmidt & Benović, 1977
<i>Corymorpha annulata</i>	1973	SA	UN	C	Schmidt & Benović, 1977
<i>Eucheilota paradoxa</i>	1967	SA	SH	E	Schmidt & Benović, 1977
<i>Eudendrium carneum</i>	<2000	-	UN	C	UNEP-MAP-RAC/SPA, 2013
<i>Eudendrium cf. merulum</i>	1969	-	UN	C	UNEP-MAP-RAC/SPA, 2013
<i>Haliscera bigelowi</i>	2002	SA	UN	C	Benović <i>et al.</i> , 2005
<i>Trichydra pudica</i>	1973	SA	UN	C	Schmidt & Benović, 1977
Mollusca					
Bivalvia					
<i>Anadara transversa</i>	2011	NA	UN	C	Nerlović <i>et al.</i> , 2012
<i>Arcuatula senhousia</i>	2003	NA	UN	C	Crocetta, 2011
<i>Brachidontes pharaonis</i>	<2006	NA	SH	C	Hrs-Brenko & Legac, 2006
<i>Bursatella leachii</i>	<1998	NA	LM	S	De Min & Vio, 1998
<i>Crassostrea gigas</i>	<1980	NA	SH	C	Hrs-Brenko, 1982
<i>Pinctada imbricata radiata</i>	2006	NA	SH	C	Dogan & Nerlović, 2008
Gastropoda					
<i>Aplysia dactylomela</i>	2006	MA	UN	S	Turk, 2006
<i>Halgerda willeyi</i>	1988	NA	UN	C	Turk, 2000
<i>Melibe viridis</i>	2001	MA	UN	C	Despalatović <i>et al.</i> , 2002
<i>Siphonaria pectinata</i>	2003	MA	SH	E	Despalatović <i>et al.</i> , 2008
Echiura					
<i>Ochetostoma erythrogrammon</i>	1962	NA	UN	C	Saiz Salinas & Ruthensteiner, 2005
Annelida					
<i>Ficopomatus enigmaticus</i>	2006	MA	SH	S	Mikuš <i>et al.</i> , 2007
<i>Metasychis gotoi</i>	1934	-	UN	C	UNEP-MAP-RAC/SPA, 2013
<i>Neopseudocapitella brasiliensis</i>	<1985	NA	UN	E	Zavodnik <i>et al.</i> , 1985
<i>Nereis persica</i>	1983	-	UN	C	UNEP-MAP-RAC/SPA, 2013
Crustacea					
<i>Arietellus pavoninus</i>	<1998	SA	SH	C	Hure & Kršinić, 1998
<i>Amphibalanus eburneus</i>	<1968	NA	SH	E	Zavodnik & Igić, 1968
<i>Amphibalanus improvisus</i>	<1986	NA	SH	E	Igić, 1986
<i>Balanus trigonus</i>	<1968	NA	SH	E	Igić, 2007
<i>Callinectes sapidus</i>	2004	SA	UN	E	Onofri <i>et al.</i> , 2007
<i>Hemigrapsus sanguineus</i>	2001	NA	UN	C	Schubart, 2003
<i>Megabalanus tintinnabulum</i>	1947	NA	SH	E	Kolosváry, 1947
Bryozoa					
<i>Bugula fulva</i>	<1998	NA	UN	C	Hayward & McKinney, 2002
Chordata					
Pisces					
<i>Cyclopterus lumpus</i>	2004	SA	SH	C	Dulčić & Golani, 2006
<i>Elates ransonnetii</i>	2010	MA	SH	C	Dulčić <i>et al.</i> , 2010
<i>Equulites (Leiognathus) klunzingeri</i>	2000	SA	LM	C	Dulčić & Pallaoro, 2002
<i>Fistularia commersonii</i>	2006	SA	LM	E	Dulčić <i>et al.</i> , 2008
<i>Holacanthus ciliaris</i>	2011	MA	SH	C	Dulčić (unpublished data)
<i>Pagrus major</i>	2004	MA	AQ	E	Dulčić & Kraljević, 2007
<i>Pampus argenteus</i>	1896	NA	LM	C	Dulčić <i>et al.</i> , 2004
<i>Paranthias furcifer</i>	2011	MA	SH	C	Pećarević & Mikuš, 2012
<i>Siganus luridus</i>	2010	SA	LM	E	Dulčić <i>et al.</i> , 2011a
<i>Siganus rivulatus</i>	2000	SA	LM	C	Dulčić & Pallaoro, 2004
<i>Sphyræna chrysoaenia</i>	2000	SA	LM	E	Pallaoro & Dulčić, 2001

Table 2. List of introduced species which have expanded their distribution range in the Eastern Adriatic Sea (Croatia) due to climate change. Legend: Location: NA – Northern Adriatic, MA – Middle Adriatic, SA – Southern Adriatic; Species origin: AT – Atlantic, ME – Mediterranean, IP – Indo-Pacific; Status: C – casual, E – established, S – spreading.

Taxon	First record	Location	Species origin	Status	Reference
Pyrrhophyta (Dinophyceae)					
<i>Akashiwo sanguinea</i>	<2011	SA, MA	AT	S	Carić <i>et al.</i> , 2011
<i>Ceratoperidinium yeye</i>	2003	NA, MA	IP	C	Ninčević Gladan <i>et al.</i> , 2006
<i>Gymnodinium abbreviatum</i>	2006	MA	AT	C	Skejić <i>et al.</i> , 2012
<i>Gymnodinium agiliforme</i>	2006	MA	AT	C	Skejić <i>et al.</i> , 2012
<i>Gymnodinium grammaticum</i>	2006	MA	AT	C	Skejić <i>et al.</i> , 2012
<i>Gymnodinium ostenfeldii</i>	2006	MA	AT	C	Skejić <i>et al.</i> , 2012
<i>Gymnodinium uberrimum</i>	2006	MA	AT	C	Skejić <i>et al.</i> , 2012
<i>Gyrodinium ascendans</i>	2006	MA	AT	C	Skejić <i>et al.</i> , 2012
<i>Gyrodinium obtusum</i>	2006	MA	AT	C	Skejić <i>et al.</i> , 2012
<i>Gyrodinium pingue</i>	2006	MA	AT	C	Skejić <i>et al.</i> , 2012
<i>Neoceratium paradoxoides</i>	2008	SA	AT	C	Batistić <i>et al.</i> , 2012
<i>Ornithocercus splendidus</i>	2008	SA	AT	C	Batistić <i>et al.</i> , 2012
<i>Scaphodinium mirabile</i>	2002	SA	AT	S	Čalić, 2010
<i>Woloszynskia neglecta</i>	2006	MA	AT	C	Skejić <i>et al.</i> , 2012
Rhodophyta					
<i>Parviphycus antipai</i>	1997	NA	AT, IP, ME	C	Sartoni & Rossi, 1998
<i>Polysiphonia banyulensis</i>	1997	NA	ME	C	Sartoni & Rossi, 1998
Cnidaria					
Anthozoa					
<i>Alicia mirabilis</i>	1998	SA	AT, ME	C	Kružić, 2002
<i>Astroides calycularis</i>	1899	NA	AT, ME	E	Zibrowius & Grieshaber, 1977
<i>Balanophyllia (Balanophyllia) regia</i>	1998	SA	AT, ME	C	Kružić, 2002
<i>Caryophyllia (Caryophyllia) cyathus</i>	1998	SA	AT, ME	C	Kružić, 2002
<i>Cladocora debilis</i>	2002	SA	AT, ME	C	Kružić <i>et al.</i> , 2007
<i>Cladopsammia rolandi</i>	2002	SA	AT, ME	E	Kružić, 2008a
<i>Coenocyathus cylindricus</i>	1998	SA	AT, ME	C	Kružić, 2002
<i>Dendrophyllia ramea</i>	1998	SA	AT, ME	C	Kružić, 2002
<i>Gyneria annulata</i>	1998	SA	AT, ME	C	Kružić, 2002
<i>Halcampoides purpureus</i>	1998	SA	AT	C	Kružić, 2002
<i>Sphenotrochus (Sphenotrochus) andrewianus</i>	1998	SA	AT, ME	C	Kružić, 2002
Hydrozoa					
<i>Lensia subtiloides</i>	1974	SA	AT	E	Batistić & Garić, 2012
<i>Muggiaea atlantica</i>	1974	SA	AT	E	Gamulin & Kršinić, 2000
<i>Niobia dendrotentaculata</i>	2001	SA	AT	E	Batistić & Garić, 2010
Mollusca					
Bivalvia					
<i>Idas simpsoni</i>	2003	SA	AT	C	Bolotin <i>et al.</i> , 2005
Gastropoda					
<i>Desmopterus papilio</i>	1993	SA	AT	C	Batistić <i>et al.</i> , 2004
<i>Protatlanta souleyeti</i>	1994	SA	AT, IP	C	Batistić <i>et al.</i> , 2004
Sipuncula					
<i>Aspidosiphon (Akrikos) mexicanus</i>	1986	NA	AT, ME	C	Murina & Zavodnik, 1986
<i>Phascolosoma scolops</i>	<1976	NA	IP	C	Murina, 1976
Annelida					
<i>Hippolyte prideauxiana</i>	2002	NA	AT, ME	C	Kirinčić, 2006
<i>Phalacrophorus pictus</i>	<2011	SA	AT	C	Batistić & Garić, 2012
<i>Pontodora pelagica</i>	<2011	SA	AT, IP	C	Batistić & Garić, 2012
Crustacea					
<i>Peniculus fistula</i>	2008	MA	AT, IP	C	Vidjak <i>et al.</i> , 2008
Chaetognatha					
<i>Sagitta galerita</i>	<2011	SA	IP	C	Batistić & Garić, 2012
Chordata					
Tunicata					
<i>Thalia orientalis</i>	2008	SA	AT	E	Batistić <i>et al.</i> , 2009
Pisces					
<i>Alectis alexandrinus</i>	1973	MA	AT, ME	C	Dulčić, 2005
<i>Caranx crysos</i>	2008	NA	AT	C	Dulčić <i>et al.</i> , 2009b
<i>Caranx rhonchus</i>	2011	SA	AT	C	Kožul & Antolović, 2013
<i>Enchelycore anatina</i>	2010	MA	AT, ME	C	Lipej <i>et al.</i> , 2011
<i>Epinephelus aeneus</i>	1998	SA	AT, ME	S	Glamuzina <i>et al.</i> , 2000
<i>Lagocephalus lagocephalus lagocephalus</i>	2004	SA	AT, IP	C	Dulčić & Pallaoro, 2006
<i>Lobotes surinamensis</i>	2010	MA	AT, IP, ME	E	Dulčić & Dragičević, 2011a
<i>Mycteroperca rubra</i>	2000	SA	AT, ME	S	Glamuzina <i>et al.</i> , 2002
<i>Plectorhinchus mediterraneus</i>	1993	NA	AT, ME	C	Lipej <i>et al.</i> , 1995
<i>Sphaeroides pachygaster</i>	1992	NA	AT, IP	S	Pallaoro & Jarda, 1996
<i>Sphyræna viridensis</i>	2003	SA	AT, ME	E	Kožul <i>et al.</i> , 2005

er, these two fish species were not included in Table 1 and the total number of non introduced species in Croatian waters.

Species biodiversity in the Adriatic Sea is influenced by water masses from the Mediterranean Sea. Mediterranean water enters the Adriatic basin from the Ionian Sea through the Strait of Otranto and flows to the northern part due to the northward coastal current. Therefore, Ionian Sea water masses, consisting of Levantine Intermediate Water and Ionian Surface Water, flow along the eastern coast of the basin. The volume of this flow varies, depending on climatic fluctuations that occur from the Atlantic to the southeast Mediterranean, but is generally greater during the winter (Batistić *et al.*, 2012).

Recorded changes in Ionian Sea circulation, consisting in the mixing of higher salinity Aegean/Levantine water with lower salinity water of Atlantic origin (Gačić *et al.*, 2010), could possibly explain the presence of Levantine and Atlantic species in the Eastern Adriatic Sea. This refers to Eastern Adriatic ichthyofauna in particular.

Regarding other introduced species in the Eastern Adriatic, possible vectors of introduction could be aquaculture activities and shipping. However, it should be noted that Croatian ports are mainly import ports thus ballast water volume is not significant. That could be the reason for fewer numbers of non indigenous planktonic species in the Eastern Adriatic in comparison with other taxa. In contrast, in the Northern Adriatic ports ballast water is an important vector of introduction of non indigenous species (David & Jakomin, 2003).

Since the Croatian part of the Adriatic Sea covers more than 35% of the total Croatian territory, monitoring of changes in marine ecosystems and biodiversity conservation are of great importance. Introduction of species in the Eastern Adriatic Sea is increasing due primarily to climate change, but other vectors should not be neglected. In order to monitor changes in Adriatic flora and fauna in the best possible way, the establishment of a national data base is necessary, as well as collaboration with other Adriatic and Mediterranean countries

Acknowledgements

This research was supported by the Ministry of Science, Education and Sports of the Republic of Croatia. We would like to extend our gratitude to all our colleagues researchers who have collected many valuable data about the Adriatic Sea during the past years. Special thanks to Dr. Argyro Zenetos for her support and useful comments.

References

Antolić, B., Špan, A., Žuljević, A., Nikolić, V., Grubelić, I. *et al.*, 2011a. A checklist of the benthic marine macroalgae

- from the eastern Adriatic coast: III. Rhodophyta 1: Ceramiales. *Acta Adriatica*, 52: 67-86.
- Antolić, B., Zavodnik, N., Špan, A., Žuljević, A., 2011b. Taxonomic composition, depth distribution and phytogeographic characteristics of marine benthic macroflora from Rijeka Bay (North Adriatic Sea, Croatia). *Natura Croatica*, 20, 53-95.
- Appeltans, W., Bouchet, P., Boxshall, G.A., De Broyer, C., de Voogd, N.J. *et al.* (eds) 2012. World Register of Marine Species. Accessed at <http://www.marinespecies.org> on 2013-03-06. (Accessed 21 February 2013)
- Batistić, M., 1999. *Macrozooplankton in the South Adriatic Pit*. PhD Thesis. University of Zagreb, Croatia, 186 pp. (in Croatian)
- Batistić, M., Garić, R., 2010. Changes in the hydrozoan community in the Adriatic Sea from 1993 to 2009. A possible relationship with hydroclimatic changes in the Mediterranean Sea. p. 15-15. In: *7th workshop of the Hydrozoan society, Porto Cesareo, Lecce, 10-18 September 2010*. Hydrozoan society, Lecce.
- Batistić, M., Garić, R., 2012. Newly recorded zooplankton species in the Adriatic Sea from 1993-2011: indicators of the hydroclimatic changes in the Eastern Mediterranean. p. 23-23. In: *International workshop: Molecular tools for monitoring marine invasive species, Lecce, 12-14 September 2012*. Italian National Interuniversity Consortium for Marine Sciences, Pisa.
- Batistić, M., Jasprica, N., Carić, M., Čalić, M., Kovačević, V. A. *et al.*, 2012. Biological evidence of a winter convection event in the South Adriatic: A phytoplankton maximum in the aphotic zone. *Continental Shelf Research*, 44, 57-71.
- Batistić, M., Jasprica, N., Carić, M., Lučić, D., 2007. Annual cycle of the gelatinous invertebrate zooplankton of the eastern South Adriatic coast (NE Mediterranean). *Journal of Plankton Research*, 29, 671-686.
- Batistić, M., Jasprica, N., Garić, R., 2009. Increasing dominance of two allochthonous gelatinous zooplankton species in the Adriatic Sea: A possible relationship with hydroclimatic changes. p. 19-19. In: *ASLO Aquatic Sciences Meeting 2009, Nice, 25-30 January 2009*. ASLO, Waco.
- Batistić, M., Kršinić, F., Jasprica, N., Carić, M., Viličić, D. *et al.*, 2004. Gelatinous invertebrate zooplankton of the South Adriatic: species composition and vertical distribution. *Journal of Plankton Research*, 26, 459-474.
- Batelli, C., Arko-Pijevec, M., 2003. Structure and seasonal variations of a subtidal turfdominated assemblage of the Oštro cape (Rijeka bay, Northern Adriatic Sea). p. 321-322. In: *8th Croatian Biological Congress, Zagreb, 27 September - 2 October, 2003*. Hrvatsko biološko društvo 1885, Zagreb.
- Bello, G., 1995. *Tylosurus acus imperialis* (Osteichthyes, Belontiidae) – a fish new to the Adriatic Sea. *Cahiers de Biologie Marine*, 36, 197-199.
- Benović, A., Lučić, D., Onofri, V., Batistić, M., Njire, J., 2005. Bathymetric distribution of medusae in the open waters of the middle and south Adriatic Sea during spring 2002. *Journal of Plankton Research*, 27, 79-89.
- Bianchi, C.N., Morri C., 1996. *Ficopomatus* 'reefs' in the Po River Delta (Northern Adriatic): their constructional dynamics, biology, and influences on the brackish-water biota. *Marine Ecology*, 17, 51-66.
- Boisset, F., 1986. Dos especies nuevas par el litoral mediterraneo peninsular: *Gelidiella antipai* M. Celan y *Myrionema*

- liechtensternii* Hauck. *Folia Botanica Miscellanea*, 5: 43-48.
- Bolotin, J., Hrs-Brenko, M., Tutman, P., Glavić, N., Kožul, V., Skaramuca, B. *et al.*, 2005. First record of *Idas simpsoni* (Mollusca, Bivalvia: Mytilidae) in the Adriatic Sea. *Journal of the Marine Biological Association of the United Kingdom*, 85: 977-978.
- Boudouresque, C.F., 1972. Vegetation marine de l'île de Port-Cros (Parc National) IX. – Sur *Gelidiella antipai* Marie Celan (Gélidiales). *Bulletin de la Société Phycologique de France*, 17: 1-8.
- Boudouresque, C.F., Verlaque, M., 2002. Biological pollution in the Mediterranean Sea: invasive versus introduced macrophytes. *Marine Pollution Bulletin*, 44: 32-38.
- Buljan, M., Zore-Armanda, M., 1976. Oceanographical properties of the Adriatic Sea. *Oceanography and Marine Biology: an annual review*, 14: 11-98.
- Čalić, M., 2010. *The influence of environmental factors on phytoplankton succession in Mali Ston Bay*. PhD Thesis. University of Zagreb, Croatia, 104 pp. (in Croatian).
- Cabioc'h, J., Flocc'h, J.H., Le Toquin, A., Boudouresque, C.F., Meinesz, A. *et al.*, 1992. *Guide des algues des mers d'Europe*. Delachaux et Nestlé, Paris, 272 pp.
- Carić, M., Jasprica, N., Čalić, M., Batistić, M., 2011. Phytoplankton response to high salinity and nutrient limitation in the eastern Adriatic marine lakes. *Scientia Marina*, 75: 493-505.
- Cornelius, P.F.S., 1981. Life cycle, dispersal and distribution among the Hydroida. *Porcupine Mewsletter*, 2, 47-50.
- Cossu, A., Gazale, V., Baroli, M., 1993. La flora marina della Sardegna: inventario delle alghe bentoniche. *Giornale Botanico Italiano*, 126, 651-707.
- Crocetta, F., 2011. Marine alien Mollusca in the Gulf of Trieste and neighbouring areas: a critical review and state of knowledge (updated in 2011). *Acta Adriatica*, 52, 247-260.
- Cukrov, M., Despalatović, M., Žuljević, A., Cukrov, N., 2010. First record of the introduced fouling tubeworm *Ficopomatus enigmaticus* (Fauvel, 1923) in the Eastern Adriatic Sea, Croatia. p. 483-483. In: 39th CIESM Congress, Venice, 10-14 May 2010. CIESM, Monaco.
- David, M., Gollasch, S., Cabrini, M., Perković, M., Bošnjak, D. *et al.*, 2007. Results from the first ballast water sampling study in the Mediterranean Sea - the Port of Koper study. *Marine Pollution Bulletin*, 54, 53-65.
- David, M., Jakomin, L., 2003. Ballast water threat in the North Adriatic – Approaching the risk assessment. In: 7th International Conference on Traffic Science, ICTS 2003, Nova Gorica, 06-08 November 2003. Faculty of Maritime Studies and Transport, Portorož.
- De Min, R., Vio, E., 1998. Molluschi esotici nell'Alto Adriatico. *Annals for Istrian and Mediterranean Studies*, 13, 43-54.
- Despalatović, M., Antolić, B., Grubelić, I., Žuljević, A., 2002. First record of the Indo-Pacific gastropod *Melibe fimbriata* in the Adriatic Sea. *Journal of the Marine Biological Association of the United Kingdom*, 82, 923-924.
- Despalatović, M., Grubelić, I., Nikolić, V., Dragičević, B., Dulčić, J. *et al.*, 2008. Allochthonous warm water species in the benthic communities and ichthyofauna of the eastern part of the Adriatic Sea. p. 51-57. In: *Climate warming and related changes in Mediterranean marine biota, Helgoland, 27 - 31 May 2008*. CIESM, Monaco.
- Dogan, A., Nerlović, V., 2008. On the occurrence of *Pinctada radiata* (Mollusca: Bivalvia: Pteriidae), an alien species in Croatian waters. *Acta Adriatica*, 49, 155-158.
- Dulčić, J., 2002. Northernmost occurrence of *Spherooides pachygaster* (Tetraodontidae) in the Adriatic Sea. *Bulletin of Marine Science*, 70, 133-139.
- Dulčić, J., 2005. On the record of the African threadfish *Alectis alexandrinus* (Pisces: Carangidae) from the Adriatic Sea. *Journal of the Marine Biological Association of the United Kingdom*, 85, 1013-1014.
- Dulčić, J., Dragičević, B., 2011a. First record of the Atlantic tripletail, *Lobotes surinamensis* (Bloch, 1790), in the Adriatic Sea. *Journal of applied ichthyology*, 27, 1385-1386.
- Dulčić, J., Dragičević, B., 2011b. *Nove ribe Jadranskog i Sredozemnog mora*. Institut za oceanografiju i ribarstvo, Split; Državni zavod za zaštitu okoliša, Zagreb, 160 pp.
- Dulčić, J., Dragičević, B., Grgičević, R., Lipej, L., 2011a. First substantiated record of a Lessepsian migrant - the dusky spinefoot, *Siganus luridus* (Actinopterygii: Perciformes: Siganidae, in the Adriatic Sea. *Acta Ichthyologica et Piscatoria*, 41, 141-143.
- Dulčić, J., Golani, D., 2006. First record of *Cyclopterus lumpus* L., 1758 (Osteichthyes: Cyclopteridae) in the Mediterranean Sea. *Journal of Fish Biology*, 69, 300-303.
- Dulčić, J., Jardas, I., Pallaoro, A., Lipej, L., 2004. On the validity of the record of silver pomfret *Pampus argenteus* (Stromateidae) from the Adriatic Sea. *Cybium*, 28, 69-71.
- Dulčić, J., Kovačić, M., Dragičević, B., 2009a. Range extension and additional records of the yellowmouth barracuda, *Sphyræna viridensis* (Actinopterygii: Perciformes: Sphyrænidae) in the eastern Adriatic Sea. *Acta Ichthyologica et Piscatoria*, 39, 59-61.
- Dulčić, J., Kraljević, M., 2007. On the record of red seabream, *Pagrus major* (Temminck and Schlegel, 1843) (Osteichthyes: Sparidae) in the Adriatic Sea. *Scientia Marina*, 71, 15-17.
- Dulčić, J., Pallaoro, A., 2002. First record of the lessepsian migrant *Leiognathus klunzingeri* (Pisces: Leiognathidae) from the Adriatic Sea. *Journal of the Marine Biological Association of the United Kingdom*, 82, 523-524.
- Dulčić, J., Pallaoro, A., 2003. First record of the filefish, *Stephanolepis diaspro* (Monacanthidae), in the Adriatic Sea. *Cybium*, 27, 321-322.
- Dulčić, J., Pallaoro, A., 2004. First record of the marbled spinefoot, *Siganus rivulatus* (Pisces, Siganidae) in the Adriatic Sea. *Journal of the Marine Biological Association of the United Kingdom*, 84, 1087-1088.
- Dulčić, J., Pallaoro, A., 2006. First record of the oceanic puffer (*Lagocephalus lagocephalus lagocephalus* Linnaeus, 1758), for the Adriatic Sea. *Journal of Applied Ichthyology*, 22, 94-95.
- Dulčić, J., Pallaoro, A., Dragičević, B., 2009b. First record of the blue runner, *Caranx crysos* (Mitchill, 1815), for the Adriatic Sea. *Journal of Applied Ichthyology*, 25, 481-462.
- Dulčić, J., Pallaoro, A., Dragičević, B., Stagličić-Radica, N., 2010. On the record of dwarf flathead *Elates ransonnetii* (Platycephalidae) in the Adriatic Sea. *Cybium*, 34, 222-223.
- Dulčić, J., Scordella, G., Guidetti, P., 2008. On the record of the Lessepsian migrant *Fistularia commersonii* (Rüppell, 1835) from the Adriatic Sea. *Journal of Applied Ichthyology*, 24, 101-102.
- Dulčić, J., Tutman, P., Čaleta, M., 2006. Northernmost occur-

- rence of the white grouper, *Epinephelus aeneus* (Perciformes: Serranidae), in the Mediterranean area. *Acta Ichthyologica et Piscatoria*, 36, 73-75.
- Dulčić J., Tutman P., Matic-Skoko S., Glamuzina B., 2011b. Six years from first record to population establishment: the case of the blue crab, *Callinectes sapidus* Rathbun, 1896 (Brachyura, Portunidae) in the Neretva River delta (South-eastern Adriatic Sea, Croatia). *Crustaceana*, 84, 1211-1220.
- Ercegović, A., 1980. Étude comparative de la végétation des basses eaux et de celle des eaux profondes de l'Adriatique centrale. *Acta Adriatica*, 21, 11-40.
- Gačić, M., Cardin, V., Eusebi Borzelli, G.L., Civitarese, G., Cardin, V. *et al.*, 2010. Can internal processes sustain reversals of the ocean upper circulation? The Ionian Sea example. *Geophysical Research Letters*, 37, 1-5.
- Gačić, M., Civitarese, G., Miserocchi, S., Cardin, V., Crise, A. *et al.*, 2002. The open-ocean convection in the Southern Adriatic: a controlling mechanism of the spring phytoplankton bloom. *Continental Shelf Research*, 22, 1897-1908.
- Galil, B.S., Clark, P.F., Carlton, J.T. (Eds.), 2011. *In the Wrong Place - Alien Marine Crustaceans: Distribution, Biology and Impacts (Invading Nature - Springer Series in Invasion Ecology, Vol. 6)*. Springer, Dordrecht, 716 pp.
- Gamulin, T., Kršinić, F., 2000. Calyophores (Siphonophora, Calyophorae) of the Adriatic and Mediterranean Seas. *Natura Croatica*, 9, 1-198.
- Giaccone, G., 1978. Revisione della flora marina de Mare Adriatico. *Annuario Parco Marino Miramare*, 6, 1-118.
- Glamuzina B., Tutman P., Geffen, J.A., Kožul, V., Skaramuca B., 2000. Premier signalement du mérrou blanc, *Epinephelus aeneus* (Serranidae) dan le mid-est de l' Adriatique. *Cybium*, 24, 306-308.
- Glamuzina, B., Tutman, P., Kožul, V., Glavić, N., Skaramuca, B., 2002. The first recorded occurrence of mottled grouper, *Mycteroperca rubra* (Serranidae), in the southeastern Adriatic Sea. *Cybium*, 26: 156-158.
- Gómez, F., Abboud-Abi, S., 2003. Records of *Ceratoperidinium* Margalef (Dinophyceae) from the Mediterranean Sea. *Vie Milieu*, 53: 43-46.
- Gómez, F., Gorsky, G., 2003. Microplankton annual cycles in the Bay of Villefranche, Ligurian Sea, NW Mediterranean. *Journal of Plankton Research*, 25, 323-339.
- Grubelić, I., Antolić, B., Despalatović, M., Grbec, B., Beg Paklar, G., 2004. Effect of climatic fluctuations on the distribution of warm-water coral *Astroides calycularis* in the Adriatic Sea: new records and review. *Journal of the Marine Biological Association of the United Kingdom*, 84, 599-602.
- Hayward P.J., McKinney, F.K., 2002. Northern Adriatic Bryozoa from the vicinity of Rovinj, Croatia. *Bulletin of the American Museum of Natural History*, 270, 139 pp.
- Hrs-Brenko, M., 1982. *Ostrea edulis* (Linnaeus) and *Crassostrea gigas* (Thunberg) larvae in the plankton of Limski kanal in the northern Adriatic Sea. *Acta Adriatica*, 23, 399-407.
- Hrs-Brenko, M., Legac, M., 2006. Inter- and intra-species relationships of sessile bivalves on the Eastern coast of the Adriatic Sea. *Natura Croatica*, 15, 203-230.
- Huber M., 2010. *Compendium of bivalves – a full-color guide to 3,300 of the world's marine bivalves – a status on Bivalvia after 250 years of research*. ConchBooks, Hackenheim, 901 pp.
- Hure, J., Kršinić, F., 1998. Planktonic copepods of the Adriatic Sea. Spatial and temporal distribution. *Natura Croatica*, 7, 1-135.
- Igić, Lj., 2007. Cirripedia of Adriatic. *Studia Marina*, 24, 1-168.
- Igić, Lj., 1986. Biološka valorizacija zaljeva Raša. Obraštaj kao biološka kontrola kod uzgoja školjkaša. *Pomorski zbornik*, 24, 555-566.
- Iveša, Lj., Jaklin, A., Devescovi, M., 2006. Vegetation patterns and spontaneous regression of *Caulerpa taxifolia* (Vahl) C. Agardh in Malinska (Northern Adriatic, Croatia). *Aquatic Botany*, 85, 324-330.
- Joksimović, A., Dragičević, B., Dulčić, J., 2008. Additional record of *Fistularia commersonii* from the Adriatic Sea (Montenegrin coast). *Marine Biodiversity Records*, 6232, 1-4.
- Kirinčić, M., 2006. First record of *Hippolyte prideauxiana* Leach, 1817 (Crustacea, Decapoda, Caridea) in the Adriatic Sea. *Acta Adriatica*, 47, 85-88.
- Kolosváry, G. 1947. Die Balaniden der Adria. *Annales Historico Naturales Musei Nationalis Hungarici*, 39, 1-88.
- Kožul, V., Antolović N., 2013. Occurrence of the false scad, *Caranx rhonchus* Geoffroy Saint-Hilaire, 1817 in the Adriatic Sea. *Journal of Applied Ichthyology*. DOI: 10.1111/jai.12086.
- Kožul, V., Tutman, P., Glavić, N., Skaramuca, B., Bolotin, J., 2005. First record of the yellowmouth barracuda, *Sphyraena viridensis* (Cuvier, 1829) (Pisces, Sphyraneidae) from Adriatic sea. *Cybium*, 29: 201-202.
- Kovalev, A.V., 2006. On the problem of Lessepsian migrations of zooplanktonic organisms. *Mediterranean Marine Science*, 7, 67-71.
- Kršinić, F., 1980. Qualitative and quantitative investigations of the tintinnids along the eastern coast of the Adriatic. *Acta Adriatica*, 21, 19-104.
- Kršinić, F. Grbec, B., 2006. Horizontal distribution of tintinnids in the open waters of the South Adriatic (Eastern Mediterranean), *Scientia Marina*, 70, 77-88.
- Kršinić, F., Njire, J., 2001. An invasion by *Muggiaea atlantica* Cunningham, 1982 in the northern Adriatic Sea in the summer of 1997 and the fate of small copepods. *Acta Adriatica* 41, 49-59.
- Kružić, P., 2002. Marine fauna of the Mljet National Park (Adriatic Sea, Croatia). I. Anthozoa. *Natura Croatica*, 11: 265-292.
- Kružić, P., 2008a. First records of *Cladopsammia rolandi* (Cnidaria: Anthozoa) in the Adriatic Sea. *Natura Croatica* 17, 9-14.
- Kružić, P., 2008b. Variations in *Posidonia oceanica* meadow structure along the coast of the Dugi Otok Island (eastern Adriatic Sea). *Journal of the Marine Biological Association of the United Kingdom*, 88, 883-892.
- Kružić, P., Radić, I., Požar-Domac, A., 2007. First record of *Cladocora debilis* (Cnidaria: Anthozoa) in the Adriatic Sea. *Journal of the Marine Biological Association of the UK*, 85: 1555-1555.
- Kružić, P., Zibrowius, H., Požar-Domac, A., 2002. Actiniaria and Scleractinia (Cnidaria, Anthozoa) from the Adriatic Sea (Croatia): first records, confirmed occurrences and significant range extensions of certain species. *Italian Journal of Zoology*, 69, 345-353.
- Lipej, L., Furlan, B., Antolović, N., Golani, D., Dulčić, J., 2011. The first record of fangtooth moray *Enchelycore*

- anatina* (Lowe, 1839) in the Adriatic Sea. *Journal of Applied Ichthyology*, 27: 1385-1386.
- Lipej, L., Spoto, M., Dulčić, J., 1996. *Plectorhinchus mediterraneus* from off north east Italy and Slovenia - the first records of the fish of the family Haemulidae from the Adriatic Sea. *Journal of fish biology*, 48, 805-806.
- Lipej, L., Mavrič, B., Orlando-Bonaca, M., Malej, A., 2012. State of the Art of the Marine Non-Indigenous Flora and Fauna in Slovenia. *Mediterranean Marine Science*, 13: 243-249.
- Lučić, D., Benović, A., Onofri, I., Batistić, M., Gangai, B. *et al.*, 2009. Planktonic cnidarians in the open southern Adriatic Sea: A comparison of historical and recent data. *Annales series historia naturalis*, 19, 27-38.
- Margalef, R., 1969. Composición específica del fitoplancton de la costa catalano-levantina (Mediterráneo occidental) en 1962-1967 (Specific structure of phytoplankton from Catalan coast (East Mediterranean). *Investigación Pesquera*, 33, 345-380.
- Meinesz, A., Belsler, T., Thibaut, T., Antolic, B., Mustapha, K.B. *et al.*, 2001. The introduced green alga *Caulerpa taxifolia* continues to spread in the Mediterranean. *Biological Invasions*, 3, 201-210.
- Mikuš, J., Pečarić, M., Bratoš Cetinić, A., 2007. WGITMO National Report Croatia 2006. p. 56-58. In: *Working Group on Introductions and Transfers of Marine Organisms (WGITMO)*, Dubrovnik, 21-23 March 2007. ICES, Copenhagen.
- Millard, N.A.H., 1975. Monograph on the Hydroida of southern Africa. *Annals of the South African Museum*, 68, 1-513.
- Monti, M., Minocci, M., Beran, A., Iveša, Lj., 2007. First record of *Ostreopsis* cf. *ovata* on macroalgae in the Northern Adriatic Sea. *Marine Pollution Bulletin*, 54, 598-601.
- Murina, G.V.V., 1976. On species composition and distribution of sipunculids in the Mediterranean Sea. Ekspedicionnye issledovanija v Sredizemnon more 74 I 75 -jrejsy Nis "Akademik A. Kovalevskij". *Naukova Dumka*, 34-39. (In Russian).
- Murina, G.V.V., Zavodnik, D., 1986. Sipuncula of the Adriatic Sea. *Thalassia Jugoslavica*, 21-22, 23-73.
- Nerlović, V., Doğan, A., Perić, L., 2012. First record of *Anadara transversa* (Mollusca: Bivalvia: Arcidae) in Croatian waters (Adriatic Sea). *Acta Adriatica*, 53, 139-144.
- Nikolić, V., Žuljević, A., Antolić, B., Despalatović, M., Cvitković, I., 2010. Rasprostranjenost invazivne crvene alge *Womersleyella setacea* (Hollenberg) R.E. Norris (Rhodophyta, Ceramiales) u Jadranskom moru. *Acta Adriatica*, 51, 195-202.
- Ninčević Gladan, Ž., Skejić, S., Marasović, I., Žuljević, A., 2006. First record of *Ceratoperidinium yeye* in the eastern Adriatic Sea. *Acta Adriatica*, 47, 207-210.
- Occhipinti Ambrogi, A., 2002. Current status of aquatic introductions in Italy. p. 311-324. In: *Invasive aquatic species of Europe - distribution, impacts and management*. E. Lepakoski, S. Gollasch, S. Olenin (Eds). Kluwer Academic Publishers, London.
- Onofri V., Dulčić J., Conides A., Matic-Skoko S., Glamuzina B., 2007. The occurrence of the blue crab, *Callinectes sapidus* Rathbun, 1896 (Decapoda, Brachyura, Portunidae) in the Eastern Adriatic (Croatian Coast). *Crustaceana*, 81, 403-409.
- Orlando-Bonaca, M., 2001. A survey of the introduced non-indigenous species in the northern Adriatic Sea. *Annals for Istrian and Mediterranean Studies*, 11, 149-158.
- Orlando-Bonaca, M., 2010. New records of non-indigenous species in Slovenian coastal waters. *Annales Series historia naturalis*, 20, 143-150.
- Pacios, I., Guerra-García, J.M., Baeza-Rojano, E., Cabezas, M.P., 2011. The non-native seaweed *Asparagopsis armata* supports a diverse crustacean assemblage. *Marine Environmental Research*, 71, 275-82.
- Pallaoro, A., Dulčić, J., 2001. First record of the *Sphyræna chrysotaenia* (Klunzinger, 1884) (Pisces, Sphyrænidae) from the Adriatic sea. *Journal of Fish Biology*, 59, 179-182.
- Pallaoro, A., Jardas, I., 1996. Ichthyological collection of the Institute of Oceanography and Fisheries in Split (Croatia). *Natura Croatica*, 5, 177-219.
- Pax, F., Müller, I., 1962. Die Anthozoenfauna der Adria. *Fauna et Flora Adriatica*, 3, 1-343.
- Pečarić, M., Mikuš, J., 2012. WGBOSV National Report Croatia 2012. p. 41-46. In: *ICES/IOC/IMO Working Group on Introduction of Transfers of Marine Organisms (WGBOSV)*, Lisbon, 12-14 March 2012. ICES, Copenhagen.
- Peharda, M., Egzeta-Balić, D., Vrgoč, N., Isajlović, I., Bogner, D., 2010. Description of bivalve community structure in the Croatian part of the Adriatic Sea - hydraulic dredge survey. *Acta Adriatica*, 51: 141-158.
- Pérès, J.M., Gamulin-Brida, H., 1973. *Biološka oceanografija*. Školska knjiga, Zagreb, 439 pp.
- Piazzini, L., Ceccherelli, G., Cinelli, F., 2001. Threat to macroalgal diversity: effects of the introduced green alga *Caulerpa racemosa* in the Mediterranean. *Marine Ecology Progress Series*, 210: 149-159.
- Polat, S., Koray, T., 2007. Planktonic dinoflagellates of the northern Levantine Basin, northeastern Mediterranean Sea. *European Journal of Protistology*, 43: 193-204.
- Poloniato, D., Ciriaco, S., Odorico, R., Dulčić, J., Lipej, L., 2010. First record of the dusky spinefoot *Siganus luridus* (Rüppell, 1828) in the Adriatic Sea. *Annales for Istrian and Mediterranean Studies, Series historia naturalis*, 20: 161-166.
- Raibaut, A., Combes, C., Benoit, F. 1998. Analysis of parasitic copepod species richness among Mediterranean fish. *Journal of Marine Systems*, 15, 185-206.
- Relini, G., 1968. Osservazioni preliminari sui balani della Rada di Vado Ligure (SV). *Bollettino dei Musei e degli Istituti biologici dell'Università di Genova*, 36, 185-190.
- Ribera, M.A., Boudouresque, C.F., 1995. Introduced marine plants, with special reference to macroalgae: mechanisms and impact. *Progress in Phycological Researches*, 11, 187-268.
- Saiz Salinas, J. I., Ruthensteiner, B., 2005. First Record of Ochetostoma for the Mediterranean Sea (Echiura). *Spixiana*, 28, 9-11.
- Sartoni, G., Rossi, S., 1998. New records for the benthic algal flora of the Northern Adriatic Sea. *Flora Mediterranea*, 8: 9-15.
- Schmidt, H. E., Benović, A., 1977. Notes on Hydromedusae (Cnidaria) from the Adriatic Sea. *Journal of the Marine Biological Association of the United Kingdom*, 57, 635-640.
- Schmidt, H.E., Benović, A., 1979. Cruises of the research vessel "Vila Velebita" in the Kvarner region of the Adriatic Sea. XII. Hydromedusae (Cnidaria). *Thalassia Jugoslavi-*

- ca, 15, 193-202.
- Schubart, C.D., 2003. The East Asian shore crab *Hemigrapsus sanguineus* (Brachyura: Varunidae) in the Mediterranean Sea: an independent human-mediated introduction. *Scientia Marina*, 67, 195-200.
- Skejić, S., Marasović, I., Ninčević Gladan, Ž., 2012. Phytoplankton assemblages at fish farm in Maslinova bay (Island of Brač). *Croatian Journal of Fisheries*, 70, 41-52.
- Streftaris, N., Zenetos, A., 2006. Alien marine species in the Mediterranean – the 100 ‘worst invasives’ and their impacts. *Mediterranean Marine Science*, 7, 87-118.
- Špan, A., Antolić, B., 1983. Prilog poznavanju fitobentosa Crnogorskog primorja (južni Jadran). *Studia Marina*, 13-14, 87-111.
- Špan, A., Antolić, B., Žuljević, A., 1998. The genus *Caulerpa* (Caulerpales, Chlorophyta) in Adriatic Sea. p. 584-585. In: 35th CIESM Congress, Dubrovnik, 1-5 June 1998. CIESM, Monaco.
- Turk, T., 2000. The opisthobranch mollusks (Cephalaspidea, Saccoglossa, Notaspidea, Anaspidea and Nudibranchia) of the Adriatic sea with special reference to the Slovenian coast. *Annales, Series historia naturalis*, 10: 161-172.
- Turk, T., 2006. *Aplysia dactylomela* from Croatia. *Sea Slug Forum. Australian Museum, Sydney*. <http://www.seaslugforum.net/showall/aplydact> (Accessed 28 August 2006)
- UNEP-MAP-RAC/SPA, 2013. Marine Mediterranean Invasive Alien Species (MAMIAS), <http://www.mamias.org> (Accessed on 12 February 2013)
- Vaniček, V., Juračić, M., Bajraktarević, Z., Čosović, V., 2000. Benthic Foraminiferal Assemblages in a Restricted Environment - An Example from the Mljet Lakes (Adriatic Sea, Croatia). *Geologia Croatica*, 53, 269-279.
- Verlaque, M., Durand, C., Huisman, J.M., Boudouresque, C.F., Le Parco, Y., 2003. On the identity and origin of the Mediterranean invasive *Caulerpa racemosa* (Caulerpales, Chlorophyta). *European Journal of Phycology*, 38, 325-339.
- Vidjak, O., Zorica, B., Sinovčić, G. 2008. First record of parasitic copepod *Peniculus fistula* von Nordmann, 1832 (Siphonostomatoidea: Pennellidae) from garfish *Belone belone* (Linnaeus, 1761) in the Adriatic Sea. *Cahiers de biologie marine*, 49, 209-213.
- Viličić, D., Marasović, I., Mioković, D. 2002. Checklist of phytoplankton in the eastern Adriatic Sea. *Acta Botanica Croatica*, 61, 57-91.
- Viličić, D., Vučak, Z., Škrivanić, A., Gržetić, Z., 1989. Phytoplankton blooms in the oligotrophic open South Adriatic waters. *Marine Chemistry*, 28, 89-107.
- Zavatarelli, M., Raicich, F., Artegiani, A., Bregant, D., Russo, A., 1998. Climatological biogeochemical characteristics of the Adriatic Sea. *Journal of Marine Systems*, 18, 227-263.
- Zavodnik, D., Igić, Lj., 1968. Zapažanja o obraštaju okoline Rovinja. *Thalassia Jugoslavica*, 4, 55-69.
- Zavodnik, D., Kovačić, M., 2000. Index of marine fauna in Rijeka bay (Adriatic Sea, Croatia). *Natura Croatica*, 9: 297-379.
- Zavodnik, D., Vidaković, J., Amoureux, L., 1985. Contribution to sediment macrofauna in the area of Rovinj (North Adriatic Sea). *Cahiers de Biologie Marine*, 26, 431-444.
- Zenetos, A., Gofas, S., Verlaque, M., Çinar, M., García Raso, E. et al., 2010. Alien species in the Mediterranean Sea by 2010. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part I. Spatial distribution. *Mediterranean Marine Science*, 11: 381-493.
- Zibrowius, H., Grieshaber, A., 1977. Scléractiniaires de l'Adriatique. *Téthys*, 7, 375-384.
- Zrnčić, S., Le Roux, F., Oraić, D., Šoštarić B., Berthe, F.C.J., 2001. First record of *Marteillia* sp. in mussels *Mytilus galloprovincialis* in Croatia. *Diseases of Aquatic Organisms*, 44, 143-148.
- Žuljević, A., Antolić, B., Onofri, V., 2003. First record of *Caulerpa racemosa* (Caulerpales: Chlorophyta) in the Adriatic Sea. *Journal of the Marine Biological Association of the United Kingdom*, 83: 711-712.
- Žuljević, A., Despalatović, M., Nikolić, V., 2010. *Zelene alge kaulerpe*. <http://jadran.izor.hr/kaulerpa> (Accessed 20 October 2012)